\$ 4 mm	<u> </u>	;;	1		11
1	2	3	4	5	6
• • •	(4	14	11	33	€ ¥ • #
7	. 8	9	10	11	12
4.	()	11	* *	A.3	: #
13	14	15	16	17	18
••	1:				ì
19	20	21	22	23	X Y




FIG 2




Fig. 4. Panel A. PCR amplification of α -fetoprotein cDNA from reverse transcribed total RNA from undifferentiated (U) and differentiated (D) R278.5 cells. Panel B. PCR amplification of α -chorionic gonadotropin and β -chorionic gonadotropin subunit cDNAs from reverse transcribed total RNA from differentiated R278.5 cells.




FIG 5

* 4 7 m.	i i	3)	1		
1	2	3	4	5	6
	(j	14	11	33	≤ ¥
7	8	9	10	11	12
4.	()	11	**	Áđ	: #
13	14	15	16	17	18
••	1:				i
19	20	21	22	23	х ү

FiG 1




FIG 2






FIG 3





X X	.23	22	57	50	61
	,			it	• •
81	Δt	91	ςτ	Þτ	cτ
∦.	r.A	7.5	11	()	, ,
75	π	70	6	8	
# # \$	3.8	Jį	7.5	,,,	
9	s	,	τ.	z	Ţ
ii	Ġ	7)	ii	1 3	4.11
					•






Fig. 4. Panel A. PCR amplification of α -fetoprotein cDNA from reverse transcribed total RNA from undifferentiated (U) and differentiated (D) R278.5 cells. Panel B. PCR amplification of α -chorionic gonadotropin and β -chorionic gonadotropin subunit cDNAs from reverse transcribed total RNA from differentiated R278.5 cells.